

CLAIMS

1. A compound of structural formula (I), as an activators of Histone acetyltransferases, *containing ring A derived from substituted benzoic acid moiety and ring B is substituted anilide* wherein:

5 **R1** is H, Methyl, Ethyl, n-Propyl, Isopropyl, n-butyl, t-butyl, C₈H₁₈, C₁₅H₂₆, C₁₅H₂₈, C₁₅H₃₀, C₁₅H₃₂;

R2 is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl;

R3 is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF₃, CCl₃, Cl₃, F, Cl, I, NO₂, CN;

10 **R4** is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF₃, CCl₃, Cl₃, F, Cl, I, NO₂, CN;

R5 is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF₃, CCl₃, Cl₃, F, Cl, I, NO₂;

R6 is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF₃, CCl₃, Cl₃, F, Cl, I, NO₂, CN; and

15 **R7** is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF₃, CCl₃, Cl₃, F, Cl, I, NO₂, CN.

2. A compound of structural formula (II) *for ring A of formula (I)* the accompanying drawings for inhibitor of Histone acetyltransferases, wherein:

20 **R1** is H, CH₃, Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy, O-Isopropoxy, n-butoxy, t-butoxy, C₈H₁₈, C₁₅H₂₆, C₁₅H₂₈, C₁₅H₃₀, C₁₅H₃₂;

R2 is H, CH₃, Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy, O-Isopropoxy, n-butoxy, t-butoxy, C₈H₁₈, C₁₅H₂₆, C₁₅H₂₈, C₁₅H₃₀, C₁₅H₃₂;

R3 is H, CH₃, Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy, O-Isopropoxy, n-butoxy, t-butoxy, C₈H₁₈, C₁₅H₂₆, C₁₅H₂₈, C₁₅H₃₀, C₁₅H₃₂;

R4 is H, CH₃, Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy, O-Isopropoxy, n-butoxy, t-butoxy, C₈H₁₈, C₁₅H₂₆, C₁₅H₂₈, C₁₅H₃₀, C₁₅H₃₂;

R5 is H, CH₃, Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy, O-Isopropoxy, n-butoxy, t-butoxy, C₈H₁₈, C₁₅H₂₆, C₁₅H₂₈, C₁₅H₃₀, C₁₅H₃₂;

30

R6 is H, CH₃, Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy, O-Isopropoxy, n-butoxy, t-butoxy, C₈H₁₈, C₁₅H₂₆, C₁₅H₂₈, C₁₅H₃₀, C₁₅H₃₂.

3. A process of preparing compounds as described in formula I or formula II by known methods.

- 5 4. A method of treating a patient suffering from diseases due defects in gene regulation predominantly or at risk of, cancer, which comprises administering to the patient a therapeutically effective amount of a compound of formula (I) to activate histone acetyltransferases or formula (II) to inhibit histone acetyltransferase or a pharmaceutically acceptable salt or solvate of these compounds.

10

15

20

25